

REMARK ON THE PAPER "ASYMPTOTIC BEHAVIOR OF POLYNOMIALS ORTHONORMAL ON A HOMOGENEOUS SET"

F. PEHERSTORFER, AND P. YUDITSKII

ABSTRACT. Minor modifications are given to prove the Main Theorem under the Blaschke (instead of Carleson) condition as well as a small historical comment.

Because of the reference [1] on our paper [4] a certain historical comment is needed.

In [4] we generalized H. Widom's Theorem [5] based on an absolutely new idea, dealing with one dimensional perturbation of a given Jacobi matrix. As it is well known if a Jacobi matrix J has the compact $E = [b_0, a_0] \setminus \cup_{j \geq 1} (a_j, b_j)$ as spectral set, then its one dimensional perturbation may have in addition to E spectral points in the gaps (one point in each gap). Since in our case, the set E is possibly a Cantor type set, an *infinite* number of spectral points X has to be added to the support of the spectral measure. A homogeneous set E possesses the following very nice property [2]. Let $B(z, z_0) = B(z, z_0; \Omega)$ be the Blaschke factor in the domain $\Omega = \overline{\mathbb{C}} \setminus E$ with zero at $z_0 \in \Omega$. From each interval (a_j, b_j) let us pick arbitrarily exactly one x_j . Then

$$\inf_k \prod_{j \neq k} |B(x_k, x_j)| > 0. \quad (1)$$

Note that the convergence of the product $\prod_j |B(z_0, x_j)| > 0$ (the Blaschke condition) corresponds to the so called Widom property of the domain Ω . So, the domain with a homogeneous boundary has even a better property: the more restrictive condition (1), the so called Carleson condition, holds for the given Blaschke product.

Thus to use our idea on a one dimensional perturbation, we were enforced to work with spectral measures supported on a homogeneous set E but also having, possibly an infinite set of mass points supported on an arbitrary (real) set X , satisfying (similar to (1)) the Carleson condition

$$\inf_{x \in X} \prod_{y \in X, y \neq x} |B(x, y)| > 0. \quad (2)$$

When the results of [4] were presented and the manuscript was submitted for publication, we recognized that the math community is much more interested in an infinite number of mass point than in a Cantor type spectral sets. We wrote [3] considering E just as a single interval. It became, almost immediately clear that the Carleson condition was too restrictive and a simple trick [3, (2.9)] allows to use only Blaschke condition.

Unfortunately, the first paper was finally published later than the second one (in this way we have, formally, a more fresh paper with the more restrictive condition [4, (6.1)] on the set X).

To prove the Main Theorem under the Blaschke condition one has to estimate [4, (6.9)] in the way [3, (2.9)], of course, following to the standard strategy in this paper (Lemma 4.2, Lemma 5.3, etc):

1. For a fixed $\epsilon > 0$ use the finite covering of Γ^* (see [4, p.139]),

$$\Gamma^* = \bigcup_{j=1}^{l(\epsilon)} \{\beta : \text{dist}(\beta, \beta_j) \leq \eta(\epsilon)\}. \quad (3)$$

2. For the exhaustion $\{X_N\}$ of X by finite sets [4, p.138], let M_N be the character of the Blaschke product B_N with zeros at $X \setminus X_N$. Since $B_N(0) \rightarrow 1$ and $M_N \rightarrow 1_{\Gamma^*}$ we can choose N so big that

$$1 - B_N(0) \leq \epsilon, \quad 1 - \Delta_N^{-1}(0) \leq \epsilon \quad (4)$$

(for the definition of Δ^α see [4, p.125]).

3. Having a finite number of the reproducing kernels (characters β_j) and a finite number of points ($X_N = \{\text{zeros of } B/B_N\}$), choose n due to the estimation

$$\begin{aligned} & \left| \sum_{X_N} \left\{ b^{n+1} K^{\beta_j} \left(\frac{-z'}{\phi'} \right) P_n \right\} \right| \\ & \leq (\sup_{X_N} |b|)^n \sqrt{\sum_{X_N} |P_n|^2 \sigma_l} \sqrt{\sum_{X_N} \left| K^{\beta_j} \frac{b}{\psi} \right|^2 \sigma_l}. \end{aligned} \quad (5)$$

Let us mention that the key Lemmas 1.1, 2.2, 2.4, 5.2 were proved under the Blaschke condition [4, (2.5)].

REFERENCES

1. D. Damanik, R. Killip, and B. Simon *Perturbation of orthogonal polynomials with periodic recursion coefficients*, Preprint.
2. P. W. Jones and D. E. Marshall, *Critical points of Green's function, harmonic measure, and the corona problem*, Ark. Mat. **23**, 281–314 (1985).
3. F. Peherstorfer and P. Yuditskii, *Asymptotics of orthonormal polynomials in the presence of a denumerable set of mass points*, Proc. Amer. Math. Soc. **129** (2001), 3213–3220.
4. F. Peherstorfer and P. Yuditskii, *Asymptotic behavior of polynomials orthonormal on a homogeneous set*, J. Analyse Math. **89**, 113–154 (2003).
5. H. Widom, *Extremal polynomials associated with a system of curves in the complex plane*, Adv. Math. **3** (1969), 127–232.